

# Oops!

Everybody makes mistakes!

Can you find the mistake in this training slide?

How would you correct it?

Now Viewing: Bloodborne Pathogens

Global Compliance Network

## Spills

OSHA recommends use of an EPA-registered disinfectant suitable for non-enveloped viruses (e.g., adenovirus, norovirus, poliovirus) to treat contamination/spills and to disinfect surfaces after bulk spill material has been removed.

Follow manufacturer instructions for appropriate concentration and contact time (refer to Safety Data Sheets) for the specific disinfectant.

When commercial disinfectant products are unavailable, common household bleach (if allowed) and other appropriate disinfectants may be effective alternatives. Use a 1:10 solution of bleach to water (e.g., 1 cup of bleach in 9 cups of water).



# Oops!

Everybody makes mistakes!  
Can you find the mistake in this training slide?

The slide adds the correction, but your students may choose to correct this by changing the ratio (1:9), or by changing the text to read solution instead of water.

How would you correct it?

Now Viewing: Bloodborne Pathogens

Global Compliance Network

## Spills

OSHA recommends use of an EPA-registered disinfectant suitable for non-enveloped viruses (e.g., adenovirus, norovirus, poliovirus) to treat contamination/spills and to disinfect surfaces after bulk spill material has been removed.



Follow manufacturer instructions for appropriate concentration and contact time (refer to Safety Data Sheets) for the specific disinfectant.

When commercial disinfectant products are unavailable, common household bleach (if allowed) and other appropriate disinfectants may be effective alternatives. Use a 1:10 solution of bleach to water (e.g., 1 cup of bleach in 9 cups of water).

The above ratio is a Part-to-Whole ratio (1 cup of bleach within the total/10 cups of the final mixture), not a Part-to-Part ratio.

PREV SLIDE    RELOAD SLIDE    ||    NEXT SLIDE

SEND A COMMENT    REQUEST ASSISTANCE    [CC] DISPLAY SLIDE TEXT